

# Addiction Severity Index 5th Edition

Clinical/Training Version

**A. Thomas McLellan, Ph.D.  
Deni Carise, Ph.D.  
Thomas H. Coyne, Ed.D., LCSW**

**Remember:**

- This is an interview, not a test.
- Items Circled are to be asked at follow-up.
- Items with an asterisk are cumulative and should be re-phrased at follow-up.

**INTRODUCING THE ASI:**

Introduce and explain the seven potential problem areas: Medical, Employment/Support Status, Alcohol, Drug, Legal, Family/Social, and Psychiatric.

All clients receive this same standard interview.

All information gathered is confidential; explain what that means in your facility; who has access to the information and the process for the release of information.

There are two time periods we will discuss:

1. The past 30 days
2. Lifetime

**Patient Rating Scale:**

Patient input is important. For each area, I will ask you to use this scale to let me know how troubled or bothered you have been by any problems in each section. I will also ask you how important treatment is for the area being discussed.

**The scale is:**

- 0 - Not at all
- 1 - Slightly
- 2 - Moderately
- 3 - Considerably
- 4 - Extremely

Inform the client that he/she has the right to refuse to answer any question. If the client is uncomfortable or feels it is too personal or painful to give an answer, instruct the client not to answer.

Explain the benefits and advantages of answering as many questions as accurately as possible in terms of developing a comprehensive and effective treatment plan to help them.

**Please try not to provide inaccurate information!**

**INTERVIEWER INSTRUCTIONS:**

1. Leave no blanks.
2. Make plenty of Comments (if another person reads this ASI, they should have a relatively complete picture of the client's perceptions of his/her problems).
3. X = Question not answered.  
N = Question not applicable.
4. Terminate interview if client misrepresents two or more sections.
5. When noting comments, please write the question number.
6. Tutorial/clarification notes are preceded with "•".

**HALF TIME RULE:**

If a question asks the number of months, round up periods of 14 days or more to 1 month. Round up 6 months or more to 1 year.

**CONFIDENCE RATINGS:**

- ⇒ Last two items in each section.
- ⇒ Do not over-interpret.
- ⇒ Denial does not warrant misrepresentation.
- ⇒ Misrepresentation = overt contradiction in information.

***Probe, cross-check and make plenty of comments!***

**HOLLINGSHEAD CATEGORIES:**

1. Higher execs, major professionals, owners of large businesses.
2. Business managers of medium sized businesses, lesser professions, i.e., nurse, opticians, pharmacists, social workers, and teachers.
3. Administrative personnel, managers, minor professionals, owners/proprietors of small businesses, i.e., bakery, car dealership, engraving business, plumbing business, florist, decorator, actor, reporter, travel agent.
4. Clerical and sales, technicians, small businesses (bank teller, bookkeeper, clerk, draftsman, timekeeper, secretary).
5. Skilled manual - usually having had training (baker, barber, brake person, chef, electrician, fireman, machinist, mechanic, paperhanger, painter, repairperson, tailor, welder, police, plumber).
6. Semi-skilled (hospital aide, painter, bartender, bus driver, cutter, cook, drill press, garage guard, checker, waiter, spot welder, machine operator).
7. Unskilled (attendant, janitor, construction helper, unspecified labor, porter, including unemployed).
8. Homemaker.
9. Student, disabled, no occupation.

**LIST OF COMMONLY USED DRUGS:**

Alcohol:	Beer, wine, liquor
Methadone:	Dolophine, LAAM
Opiates:	Pain killers = Morphine, Diluaid, Vicodin, Oxycontin, Hydrocodone, Demerol, Percocet, Darvon, Talwin, Codeine, Tylenol 2,3,4, Robitussin, Fentanyl
Barbiturates:	Nembutal, Seconal, Tuinol, Amytal, Pentobarbital, Secobarbital, Phenobarbital, Fiorinol
Sed/Hyp/Tranq:	Benzodiazepines = Valium, Librium, Ativan, Serax Tranxene, Xanax, Miltown, Other = ChloralHydrate (Noctex), Quaaludes Dalmane, Halcion
Cocaine:	Cocaine Crystal, Free-Base Cocaine or "Crack, and "Rock Cocaine"
Amphetamines:	Monster, Crank, Benzedrine, Dexedrine, Ritalin, Preludin, Methamphetamine, Speed, Ice, Crystal
Cannabis:	Marijuana, Hashish
Hallucinogens:	LSD (Acid), Mescaline, Mushrooms (Psilocybin), Peyote, Green, PCP (Phencyclidine), Angel Dust, Ecstasy
Inhalants:	Nitrous Oxide, Amyl Nitrate (Whippits, Poppers), Glue, Solvents, Gasoline, Toluene, Etc.
Just note if these are used:	Antidepressants, Ulcer Meds = Zantac, Tagamet Asthma Meds = Ventoline Inhaler, Theodur Other Meds = Antipsychotics, Lithium

**ALCOHOL/DRUG USE INSTRUCTIONS:**

The following questions refer to two time periods: the past 30 days and lifetime.

- 30 day questions only require the number of days used.
- Lifetime refers to the time prior to the last 30 days.
- Lifetime use is asked to determine extended periods of use.

**Regular Use** = 3+ times per week, binges, or problematic irregular use in which normal activities are compromised.

**Alcohol to Intoxication** does not necessarily mean "drunk", use the words "to feel or felt the effects", "got a buzz", "high", etc. instead of intoxication. As a rule, 3+ drinks in one sitting, or 5+ drinks in one day defines "intoxication".

**How to ask these questions:**

- "How many days in the past 30 have you used....?"
- "How many years in your life have you regularly used....?"























