

**ASITRAINING.com**  
**On Line Resource Center for the Addiction Severity Index**

**Follow Up ASI**  
**Clinical Training Version**

A. Thomas McLellan, Ph.D.  
Deni Carise, Ph.D.  
Laura MacPherson  
Thomas H. Coyne, MSW

**Remember: This is a follow up interview, not a test**  
**Questions will be reworded depending on the follow up time frame being used.**

**3mos. Post Intake: How many times in the last three months ....**

**6 mos. Post Intake: How many times in the past six months .....**

**Post Discharge: How many times since your discharge.....**

**INTRODUCING THE ASI:** Seven potential problem areas: Medical, Employment/Support Status, Alcohol, Drug, Legal, Family/Social, and Psychological. All clients receive this same standard interview. All information gathered is confidential.

There are two time periods we will discuss:

1. The past 30 days
2. The time that has passed since the baseline interview

**Patient Rating Scale:** Patient input is important. For each area, I will ask you to use this scale to let me know how bothered you have been by any problems in each section. I will also ask you how important treatment is for you for the area being discussed.

The scale is:

- 0 - Not at all
- 1 - Slightly
- 2 - Moderately
- 3 - Considerably
- 4 - Extremely

If you are uncomfortable giving an answer, then don't answer.

**Please do not give inaccurate information!**

**INTERVIEWER INSTRUCTIONS:**

1. Leave no blanks.
2. Make plenty of Comments (if another person reads this ASI, they should have a relatively complete picture of the client's perceptions of his/her problems).
3. X = Question not answered.  
N = Question not applicable.
4. Terminate interview if client misrepresents two or more sections.
5. When noting comments, please write the question number.
6. Tutorial/clarification notes are preceded with "•".

**HALF TIME RULE:** If a question asks the number of months, round up periods of 14 days or more to 1 month. Round up 6 months or more to 1 year.

**CONFIDENCE RATINGS:** ⇒ Last two items in each section.  
⇒ Do not over interpret.  
⇒ Denial does not warrant misrepresentation.  
⇒ Misrepresentation = overt contradiction in information.

**Probe and make plenty of comments!**

**HOLLINGSHEAD CATEGORIES:**

1. Higher execs, major professionals, owners of large businesses.
2. Business managers of medium sized businesses, lesser professions, i.e., nurses, opticians, pharmacists, social workers, teachers.
3. Administrative personnel, managers, minor professionals, owners/proprietors of small businesses, i.e., bakery, car dealership, engraving business, plumbing business, florist, decorator, actor, reporter, travel agent.
4. Clerical and sales, technicians, small businesses (bank teller, bookkeeper, clerk, draftsman, timekeeper, secretary).
5. Skilled manual - usually having had training (baker, barber, brakeman, chef, electrician, fireman, lineman, machinist, mechanic, paperhanger, painter, repairman, tailor, welder, policeman, plumber).
6. Semi-skilled (hospital aide, painter, bartender, bus driver, cutter, cook, drill press, garage guard, checker, waiter, spot welder, machine operator).
7. Unskilled (attendant, janitor, construction helper, unspecified labor, porter, including unemployed).
8. Homemaker.
9. Student, disabled, no occupation.

**LIST OF COMMONLY USED DRUGS:**

Alcohol:	Beer, wine, liquor
Methadone:	Dolophine, LAAM
Opiates:	Pain killers = Morphine, Dilaudid, Demerol, Percocet, Darvon, Talwin, Codeine, Tylenol 2,3,4, Syrups = Robitussin, Fentanyl
Barbiturates:	Nembutal, Seconal, Tuinol, Amytal, Pentobarbital, Secobarbital, Phenobarbital, Fiorinol
Sed/Hyp/Tranq:	Benzodiazepines = Valium, Librium, Ativan, Serax Tranxene, Dalmane, Halcion, Xanax, Miltown, Other = ChloralHydrate (Noctex), Quaaludes
Cocaine:	Cocaine Crystal, Free-Base Cocaine or "Crack, and "Rock Cocaine"
Amphetamines:	Monster, Crank, Benzedrine, Dexedrine, Ritalin, Preludin, Methamphetamine, Speed, Ice, Crystal
Cannabis:	Marijuana, Hashish
Hallucinogens:	LSD (Acid), Mescaline, Mushrooms (Psilocybin), Peyote, Green,
Inhalants:	PCP (Phencyclidine), Angel Dust, Ecstasy Nitrous Oxide, Amyl Nitrate (Whippits, Poppers), Glue, Solvents, Gasoline, Toluene, Etc.

Just note if these are used:

- Antidepressants,
- Ulcer Meds = Zantac, Tagamet
- Asthma Meds = Ventoline Inhaler, Theodur
- Other Meds = Antipsychotics, Lithium

**ALCOHOL/DRUG USE INSTRUCTIONS:**

The following questions refer to two time periods: the past 30 days and lifetime. Lifetime refers to the time prior to the last 30 days.

- ⇒ 30 day questions only require the number of days used.
- ⇒ Lifetime use is asked to determine extended periods of use.
- ⇒ Regular use = 3+ times per week, binges, or problematic irregular use in which normal activities are compromised.
- ⇒ Alcohol to intoxication does not necessarily mean "drunk", use the words "felt the effects", "got a buzz", "high", etc. instead of Intoxication. As a rule 3+ or 5+ drinks in one setting, or within a brief period of time defines "intoxication".
- ⇒ How to ask these questions:  
→ "How many days in the past 30 have you used...?"  
→ "How many years in your life have you regularly used...?"





















